

February 2023

420 Cooper Street, Ottawa, ON K2P 2N6

613.233.4443 | TTY: 613.233.0651 | Fax: 613.233.3987

www.centretownchc.org | info@centretownchc.org







Guide for Self-Administration of **Subcutaneous Injections**



Following these steps can help you feel more confident to safely inject your medication. You can do it!

Step 1: Control your environment

 Find a quiet, clean area that will allow you to focus and be free of distractions

Step 2: Know your rights!

- ⇒ Review the 5 medication rights before every injection:
 - Right person
 - Right medication
 - Right dose
 - Right route
 - Right time

Inspect the medication vial:

- Expiry date
- Colour, no floaters





Stay organized!

Use a cellphone or a calendar to remind yourself of your **injection date** and **injection site**. Setting up an alarm can also help you remember.




Tips and Tricks!

- If you are experiencing anxiety that is making it difficult to give yourself your injection, you may want to try one of the following:
 - Distraction: put on your favourite song or a tv show
 - Support: have a support person sit with you during your injection
 - Reward: reward yourself after your injection







Where to buy injection supplies:


- Swift Compounding Pharmacy: 276 Bank St., Ottawa (613) 422-2202
- Ontario Medical Supply (OMS): 1100 Algoma Rd. Ottawa 613-244-8620
- You may also be able to buy injection supplies at your local pharmacy.


Step 6: The injection

- Clean the skin in a circular motion with an alcohol swab. Let dry for 10 seconds.
- Remove the needle protector from your injection needle.

Subcutaneous:


- Gently pinch the skin & fatty tissue between thumb & index finger with non-dominant hand
- Insert needle in a steady motion and at a 45° or 90° angle
- Push the medication in and wait 10 seconds after injecting before withdrawing the needle. Note: medication is thick, so it can be hard to push the plunger.
- Withdraw needle and let go of your skin.
- Apply band aid if needed
- Dispose of needle in sharps container.







Pinch and inject




Step 3: Wash your hands

- Wash your hands with soap & water for 30 seconds
- Dry your hands

Step 4: Prepare your equipment

⇒ Gather what you need:

- 1. Medication vial
- 2. Alcohol swabs
- 3. Syringe: 1cc or 3cc depending on preference/dose
- 4. Drawing needle: 18G 22G
- 5. Injecting needle:
 - Subcutaneous: 25G 5/8 inch needle
- 6. Band aid and cotton ball
- 7. Sharps disposal box






Wondering where to purchase supplies? See last page →




Step 4: Draw up the medication

⇒ Prepare to draw:



 Open syringe & needle packages while keeping the tips sterile.




• Remove needle protector.

⇒ Draw the medication:

- Get some air in the syringe by pulling back on the plunger, equal to the amount of medication you will be injecting.
- Holding the vial steady on your clean surface, insert the needle by puncturing the rubber stopper of the vial, then push the air in.
- Flip the vial upside down while keeping the needle tip submerged, and pull back on plunger to withdraw the medication to your prescribed dose.



Tip: You want to avoid large air bubbles in your syringe. You can draw up more medication and push it back in, or gently tap the syringe to remove the bubbles.




- Remove drawing needle and place in sharps container.
- Attach the injection needle and gently push the medication up to the hub of the needle.

Step 5: Find your spot:

Subcutaneous

⇒ Abdomen


 Your injection site is in the fatty tissue of your abdomen, at least 1 inch away from your belly button, and anywhere between the bottom of your ribs & above your hips.
You can inject while sitting down



⇒ Upper thigh

 Your injection site is in the fatty tissue in the in the middle to outer portion of your upper thigh. You can inject while sitting down







Remember:

- Always inspect the site prior to injecting. Avoid scar tissue. Watch for lumps, redness or swelling, bruises or irritation.
- Always **rotate sites** with each injection.