



February 2023

## Guide for Self-Administration of Subcutaneous Injections

420 Cooper Street, Ottawa, ON K2P 2N6

613.233.4443 | TTY : 613.233.0651 | Fax : 613.233.3987

[www.centretownchc.org](http://www.centretownchc.org) |



@CentretownCHC



facebook.com/CentretownCHC



Following these steps can help you feel more confident to safely inject your medication. You can do it!

### Step 1: Control your environment

- Find a quiet, clean area that will allow you to focus and be free of distractions

### Step 2: Know your rights!

⇒ **Review the 5 medication rights before every injection:**

- Right person
- Right medication
- Right dose
- Right route
- Right time



#### Stay organized!

Use a cellphone or a calendar to remind yourself of your **injection date** and **injection site**. Setting up an alarm can also help you remember.




### Tips and Tricks!

- If you are experiencing anxiety that is making it difficult to give yourself your injection, you may want to try one of the following:
  - Distraction: put on your favourite song or a tv show
  - Support: have a support person sit with you during your injection
  - Reward: reward yourself after your injection

### Inspect the medication vial:

- Expiry date



- Colour, no floaters

### Where to buy injection supplies:

- Swift Compounding Pharmacy: 276 Bank St., Ottawa (613) 422-2202
- Ontario Medical Supply (OMS): 1100 Algoma Rd. Ottawa 613-244-8620
- You may also be able to buy injection supplies at

your local pharmacy.




## Step 6: The injection

- Clean the skin in a circular motion with an alcohol swab. Let dry for 10 seconds.
- Remove the needle protector from your injection needle.

### Subcutaneous:


- Gently pinch the skin & fatty tissue between thumb & index finger with non-dominant hand
- Insert needle in a steady motion and at a 45° or 90° angle
- Push the medication in and wait 10 seconds after injecting before withdrawing the needle. Note: medication is thick, so it can be hard to push the plunger.
- Withdraw needle and let go of your skin.
- Apply band aid if needed
- Dispose of needle in sharps container.



Subcutaneous Injection



Pinch and inject



## Step 3: Wash your hands

- Wash your hands with soap & water for 30 seconds
- Dry your hands

## Step 4: Prepare your equipment

### ⇒ Gather what you need:

1. Medication vial
2. Alcohol swabs
3. Syringe: 1cc or 3cc depending on preference/dose
4. Drawing needle: 18G - 22G
5. Injecting needle:
  - Subcutaneous: 25G 5/8 inch needle
6. Band aid and cotton ball
7. Sharps disposal box




Wondering where to purchase supplies? See last page →

## Step 4: Draw up the medication



### ⇒ Prepare to draw:

- Clean the rubber stopper of the medication vial for 10 seconds with an alcohol swab. Let dry for 20 seconds.
- Open syringe & needle packages while keeping the tips sterile.
- Connect the drawing needle to the syringe, twisting in a clockwise motion.
- Remove needle protector.




### ⇒ Draw the medication:

- Get some air in the syringe by pulling back on the plunger, equal to the amount of medication you will be injecting.
- Holding the vial steady on your clean surface, insert the needle by puncturing the rubber stopper of the vial, then push the air in.
- Flip the vial upside down while keeping the needle tip submerged, and pull back on plunger to withdraw the medication to your prescribed dose.



**Tip:** You want to avoid large air bubbles in your syringe. You can draw up more medication and push it back in, or gently tap the syringe to remove the bubbles.



- Remove drawing needle and place in sharps container.
- Attach the injection needle and gently push the medication up to the hub of

## Step 5: Find your spot:

### Subcutaneous

#### ⇒ Abdomen

- Your injection site is in the fatty tissue of your abdomen, at least 1 inch away from your belly button, and anywhere between the bottom of your ribs & above your hips. You can inject while sitting down

#### ⇒ Upper thigh

- Your injection site is in the fatty tissue in the middle to outer portion of your upper thigh. You can inject while sitting down

the needle.

## Remember:

- Always inspect the site prior to injecting. Avoid scar tissue. Watch for lumps, redness or swelling, bruises or irritation.
- Always **rotate sites** with each injection.